



Bioscene

Bioscene

Volume- 22 Number- 03

ISSN: 1539-2422 (P) 2055-1583 (O)

www.explorebioscene.com

Temporal Dynamics of Physico-Chemical Characteristics of the Freshwater Ecosystem in Pimpalgaon (Wakhaji) Dam, Nagpur District (M.S)

¹ Sumit S. Gurchal*;² Manisha M. Bhatkulkar;³ Jatinderpal Singh &
⁴ Manpreet Kaur Saini

¹Department of Zoology, Rashtrasant Tukadoji Maharaj Nagpur University,
Nagpur, India

²Prof. & Head, Department of Zoology, Jawaharlal Nehru Arts Commerce &
Science College, Wadi, Nagpur, India

³Department of Zoology, Baring Union Christian College, Batala (Punjab) India


⁴Centre for One Health, Guru Angad Dev Veterinary and Animal Sciences
University, Ludhiana, India

*Corresponding Author: **Sumit S. Gurchal**

Abstract: Analysis of water quality is important criterion for evaluating suitability of water for drinking and irrigation. The present study investigates the seasonal dynamics of key physio-chemical parameters in the freshwater ecosystem of Pimpalgaon (Wakhaji) dam of Nagpur District, Maharashtra, India. Water samples were collected monthly from five distinct sites of dam between June 2023 to May 2024. Thirteen standard parameters were assessed using APHA (1999, 2017) guidelines including temperature, turbidity, electrical conductivity (EC), total dissolved solids (TDS), pH, dissolved oxygen (DO), free CO₂, alkalinity, chloride, phosphate, calcium, nitrate and biological oxygen demand (BOD). Results revealed clear seasonal patterns: DO ranged from 6.72 to 7.84 mg/L, BOD varied between 2.31 to 3.1 mg/L and pH fluctuated from 6.94 to 7.72 while elevated. The interplay of these parameters significantly influences aquatic productivity, nutrient cycling and biotic composition, particularly impacting distribution of zooplankton and ichthyofaunal assemblages. This study provides baseline data essential for ecological monitoring and future sustainable water resource management strategies and mitigate anthropogenic stressors affecting freshwater reservoir ecosystems.

Keywords: Analysis, Physio-chemical Characteristics, Water Quality

Graphical Abstract



Introduction

Dams play a significant role in water storage, agriculture and electricity generation. Physico-chemical characters have a great importance as their relation with fluctuation of zooplankton and necessary for pisciculture (Ayoola and Ajani, 2009). Study of various parameters regarding dam water is necessary for understanding the metabolic events of aquatic ecosystem (Shinde et al., 2011). Aquatic biota gets depleted due to indiscriminate use of chemical fertilizers and pesticides in farming causes pollution in aquatic environment (Khan et al., 2012). Hydrological condition of water affects aquaculture activities, species composition of avifauna, eutrophication and biodiversity (Wankhade et al., 2012). Water is important natural resource for all living being, food production and economy growth. Physico-chemical characteristics also influence and regulate flora and fauna (Aazamiet al., 2015). Health of human is directly connected with quality of water and it remains as major concern for man (Dirican, 2015). Decrease in percentage of rainfall and demand of water increasing day by day as increase in population of India (Ajagekar et al., 2011). Quality of water with regulating biomass,

tropic levels, energy and biodiversity at a given time and space act as limiting factor (Wanjare and Tanpure,2020).




Figure 1. Influence of Dam Physico-Chemical Attributes on Aquatic Ecosystems, Fisheries, Biodiversity, and Human Health

Immune system of fishes affected by change in water quality directly or indirectly through pollution (Bassem, 2020). Water pH, temperature, free CO₂, alkalinity like parameters play important role in growth and development of ichthyofauna. Any variation in these characters of water affects the growth and development of fishes. Due to contamination many water-borne diseases arise which suffers population of human (Makode, 2020). Physico-chemical characteristics analysis also helps to initiate necessary affirmative actions against contamination of water bodies (Gaikwad et al., 2021). So, it is mandatory to analyse important characteristics of water quality for aquatic ecosystem. To formulate appropriate management strategies and protect rich biodiversity of dam adequate information about various parameters and delicate dynamics sustained by dam have great importance. At regular intervals, quality of water must be checked. The study aims to provide insights into its quality dynamics throughout the year to aid in environmental management and policy decisions.

Table 1:Importance of Physico-Chemical Characteristics of Dam Water

Parameter / Aspect	Importance in Aquatic Ecosystem	Impact on Biota & Human Health
Water Storage in Dams	Facilitates irrigation, potable water supply, and electrical production	Facilitates economic expansion and agricultural output
Physico-chemical Characters	Impact Fluctuations in zooplankton, aquaculture, fish well-being	Regulate plant life, animal species, and aquatic biodiversity
Chemical Fertilizers & Pesticides	Induce marine pollution	Depletion of aquatic biota
Hydrological Conditions	Impact aquaculture, eutrophication and avifaunal species composition	Impact biodiversity and ecological equilibrium
Water Quality and Rainfall	Decreased precipitation diminishes water quality amid increasing demand.	Constraints on biomass, trophic strata, energy cycles
pH, Temperature, Free CO ₂ , Alkalinity	Crucial for the growth and development of fish species	Variations influence fish immunity, reproduction, and survival.
Water Contamination	Induces waterborne illnesses in people	Direct threat to public health
Water Quality Monitoring	Helpful in identifying sources of pollution and contamination	Establishes a foundation for governance and policy formulation
Management Strategies	Consistent monitoring of water quality contributes to sustainability.	Safeguards biodiversity and guarantees safe utilization

Materials and Methods

Study Area: The present study was conducted at Pimpalgaon (Wakhaji) dam located in Narkhed Taluka, Nagpur District, Maharashtra, India. The dam is geographically positioned at latitude 21.45°N and longitude 78.55°E, approximately 3.6 km east of Narkhed. This dam is the largest irrigation reservoir in Taluka, supports local agricultural practices and biodiversity. Hence, Pimpalgaon (Wakhaji) dam plays a vital role in agriculture.




Figure 2.Location of the study area A) Map of India highlighting the state of Maharashtra; B)Maharashtra map showing Nagpur District; C)Satellite image showing location of Pimpalgaon (Wakhaji) dam (symbol of heart) within Nagpur district, Maharashtra; D) Zoomed satellite view of precise location of Pimpalgaon (Wakhaji) dam indicating general geographic region. Map A prepared from (www.MapChart.net) and B from Wikimedia Commons(File:Nagpur in Maharashtra (India).svg),licensed under CC BY-SA 3.0. for illustrative purpose while Map C and D were previously used in a study of ichthyofaunal diversity (Gurchal and Bhatkulkar,2024) and it is reused here to illustrate the same location of Pimpalgaon (Wakhaji) for physico-chemical analysis.

Sampling Design: Water samples were collected monthly from five pre-selected sampling stations in the dam for a continuous period of 12 months from June 2023 to May 2024. The stations were chosen to cover diverse ecological zones within the dam. Sampling was consistently carried out during early morning hours (7.00 A.M to 9.00 A.M) to avoid diurnal variations and maintain consistency.

Sampling Methodology: Water samples were collected from surface (approx.15-30 cm depth) using pre-cleaned polyethylene bottles of one litre capacity from five

different sites of Pimpalgaon (Wakhaji) dam. Bottles were rinsed thrice with sample water before final collection.

Parameters Analyzed: Thirteen physico-chemical parameters were analyzed:

- **Physical:** Temperature, Electrical Conductivity (EC), Total Dissolved Solids(TDS), Turbidity
- **Chemical:** pH, Alkalinity, Free Carbon Dioxide (CO₂), Dissolved Oxygen (DO), Chloride, Nitrate, Calcium, Phosphate, Biological Oxygen Demand

All physico-chemical parameters estimated as per standard procedures outlined in the APHA(1999 and 2017) manuals. To ensure data accuracy all samples were processed in triplicate and mean values with standard deviation were calculated. Instruments were calibrated before use and reagent blanks were run for chemical tests. Any deviations from standard limits were cross-checked through repeat analysis.




Figure 3.Figure depicting 13 physio-chemical parameters including physical and chemical parameters

Results

The present study was carried out to analyze and determine physico-chemical status of Pimpalgaon (Wakhaji) dam water. The data collection spanned a full annual cycle (June 2023 to May 2024). To maintain diurnal consistency all samples collected consistently between 7:00 AM to 9:00 AM morning. The monthly variation in key physico-chemical parameters (temperature, electrical conductivity, turbidity, total dissolved solids, pH, free CO₂, dissolved oxygen, chloride, nitrate,

calcium, phosphate, biological oxygen demand) including mean \pm SD is represented graphically in Figures 4 to 6 for study period of June 2023 to May 2024.





Figure 4. Graphs presented seasonwise monthly variation in key physico-chemical parameters (A-Temperature, B-Electrical Conductivity) of water in Pimpalgaon (Wakhaji) dam during study period (June 2023-May 2024). Data from five sampling stations were averaged for each parameter.






Figure 5. Graphs presented seasonwise monthly variation in key physico-chemical parameters (A-Turbidity, B-Total Dissolved Solids,C-pH,D-Free CO₂,E-Total Alkalinity,F-Chloride) of water in Pimpalgaon (Wakhaji) dam during study period (June 2023-May 2024). Data from five sampling stations were averaged for each parameter.






A) Monthly Variation in Dissolved Oxygen (mg/L)**B) Monthly Variation in Calcium (mg/L)****C) Monthly Variation in Nitrate (mg/L)****D) Monthly Variation in Phosphate (mg/L)****E) Monthly Variation in Biological Oxygen Demand (mg/L)**

Figure 6. Graphs presented seasonwise monthly variation in key physico-chemical parameters (A- Dissolved Oxygen, B-Calcium, C-Nitrate, D-Phosphate, E-Biological Oxygen Demand) of water in Pimpalgaon (Wakhaji) dam during study period (June 2023-May 2024). Data from five sampling stations were averaged for each parameter.

Temperature: Water temperature clearly exhibited a clear seasonal trend with lowest value ranging from 21.28 ± 0.164 °C to highest of 30.3 ± 0.291 °C. Monsoon season presented relatively stable temperatures, whereas winter showed a decline and summer recorded progressively higher values.

Electrical Conductivity: Electrical conductivity values lowest values ranging from 324.96 ± 2.165 umho/cm to highest of 463.54 ± 1.705 umho/cm. Electrical conductivity values showed a rising trend from June to August peaking at 463.54 umho/cm, likely due to increased surface runoff during early monsoon that brought dissolved salts and ions into the reservoir. A gradual decline in EC was observed from September onward, reaching its lowest value in May-2024, which may be attributed to dilution effects during post-monsoon and winter.

Turbidity: Turbidity of water ranged from minimum of $4.28 \text{ NTU} \pm 0.16 \text{ NTU}$ in May 2024 to a maximum of $10.34 \text{ NTU} \pm 0.421 \text{ NTU}$. The values gradually increased from June, peaked during the post-monsoon period (October-November) and declined progressively through the winter and summer months. The slight increase in turbidity observed during October coincides with peak levels of nitrate and phosphate indicating nutrient induced algal and suspended matter accumulation during this time.

Total Dissolved Solids: Total dissolved solids values highest values were during August-2023 (296.8 ± 3.67 mg/L) and lowest in summer months with recorded in May-2024 (206.48 ± 2.28 mg/L). TDS values exhibited a seasonal trend reaching a maximum of 296.8 mg/L in August 2023, which coincides with the peak monsoon period. This rise is likely due to runoff from surrounding catchment areas carrying dissolved minerals, nutrients and organic matter into the reservoir. A steady decline was observed thereafter with TDS reaching its lowest concentration of 206.48 mg/L in May 2024, likely due to reduced inflow, sedimentation of particulates and possible biological uptake in the summer months.

pH: Concentration of hydrogen ions indicated by values of pH. Present assessment shows pH range from 6.94 ± 0.05 to 7.72 ± 0.083 . Indicating the water was slightly acidic to slightly alkaline throughout the year. Influence of rainwater runoff, organic matter and reduced photosynthetic activity might tends to lower pH as lowest values were observed during monsoon (July-2023 to September-2023). pH reach peak in summer months (April-2024 to May-2024). This rise may be due to

enhanced photosynthesis by aquatic plants and algae which utilizing CO₂ and thereby raise the pH.

Free CO₂: The maximum value of 6.56 ± 0.207 mg/L was recorded in month of August-2023 and minimum value of 4.04 ± 0.114 mg/L in month of May-2024. Highest concentration observed during monsoon as likely due to increased organic matter decomposition and lower photosynthetic activity under cloudy conditions which limits CO₂ utilization.

Total Alkalinity: Alkalinity levels in dam water ranged from lowest 86.4 ± 0.604 mg/L to highest 108.6 ± 1.176 mg/L. The lowest values were observed during monsoon season (July-2023 to September-2023), possibly due to dilution caused by heavy rainfall and inflow of freshwater reduces concentration of carbonates and bicarbonates. Alkalinity peaks in summer months (March-2024 to May-2024) after gradual increase through winter. This increase may be attributed to evaporation, concentration of dissolved salts and higher biological activity releasing bicarbonates into water.

Chloride: Lowest chloride levels were recorded of 23.36 ± 0.861 mg/L in month of September-2023 due to dilution effects from monsoon rains and maximum value of 33.28 ± 0.192 mg/L in month of May-2024 because increased evaporation and concentration of salts in pre-monsoon rains.

Dissolved Oxygen: The maximum value of dissolved oxygen was 7.84 ± 0.114 mg/L recorded in February-2024 and minimum value of 6.72 ± 0.130 mg/L recorded in July-2023. Low dissolved oxygen observed in July-2023 (6.72 mg/L) and August-2023 (6.78 mg/L) which may be due to reduced photosynthetic activity due to cloud cover and increased microbial decomposition of organic matter during monsoon.

Calcium: Calcium is helpful for shell construction and also promotes growth of micro-organisms. The minimum value of calcium was 30.56 ± 0.531 mg/L recorded in September-2023 and maximum value of 42.6 ± 1.45 mg/L found in May-2024. Due to dilution by rainfall and surface runoff calcium low levels were observed in monsoon period (June-2023 to September-2024). Reduced rainfall and increased mineral concentration might lead to gradual increase from October-2023 and high concentration occurred in summer (February-2024 to May-2024).

Nitrate: Nitrate levels ranged from (12.1 ± 0.158 mg/L) in May 2023 to (25.36 ± 0.181 mg/L) of October and gradual decline toward May 2024, where values reached around 12.1 mg/L. The peak during post-monsoon suggests enhanced nutrient input likely due to agricultural runoff and organic decomposition.

Phosphate: The value of phosphate fluctuates from (15.8 ± 0.158 mg/L) to (35.36 ± 0.207) during study period. Phosphate concentration showed a trend rising

from around 17.7 mg/L in June to maximum of 35.36 mg/L in October then progressively declines through winter and summer months. The synchronous peaks of both nutrients during October highlight a strong correlation, potentially linked to increased external loading and biological activity during this period.

Biological Oxygen Demand: Biological oxygen demand fluctuated from 2.31 ± 0.089 mg/L rises till 3.1 ± 0.1 mg/L during study period. Comparatively mild high BOD levels were observed during monsoon months (July-August) likely due to increased runoff and organic input, while the low values occurred in winter months (December-February) when water temperature was lower and DO levels were high. BOD values remained within acceptable limit which indicate satisfactory water quality and minimal organic pollution pressure in dam.

Discussion

The present study assessed the monthly variations with season in physico-chemical parameters of Pimpalgaon (Wakhaji) dam of Nagpur. Water samples were collected monthly from five designated stations between June-2023 to May-2024 during morning hours (7:00-9:00).

Temperature ranged from 21.28°C (January-2024) to 30.3°C (May-2024) reflecting natural seasonal transitions from winter to summer. Summer months (March-May) recorded the highest water temperatures influencing the increase in Biological Oxygen Demand (BOD) and reduction in Dissolved Oxygen (DO) due to enhanced metabolic activities and microbial oxygen consumption. Similar seasonal temperature variations affecting Dissolved Oxygen and Biological Oxygen Demand have been reported in Anandpur Dam, Gujarat (Vyas et al., 2024).

pH values remained within a narrow and slightly alkaline range (6.94-7.72) which is favourable for zooplankton and aquatic organisms. Slight alkalinity is necessary for maintaining buffering of carbonate and promoting biological productivity. Comparable pH stability supporting zooplankton diversity was observed in Yaldali dam, Parbhani (Rathod & Walale, 2022).

The electrical conductivity and total dissolved solids values showed seasonal variation with higher levels observed during monsoon and post monsoon months (August-October) due to increased runoff and mineral input. Lower values during summer and winter indicate reduced inflow and evaporation effects. Monsoon and post monsoon months may result from runoff carrying suspended particles, increased suspended solids and organic debris affecting primary productivity by reducing light penetration. This suggests that nutrient driven turbidity possibly from algal growth and organic runoff may temporarily elevated turbidity. The observed decline in turbidity during winter and summer can be attributed to settling of sediment and decreased biological productivity. These findings are similar with previous studies that report seasonal variation in electrical conductivity, total dissolved solids and turbidity corresponding with nutrient enrichment and post-monsoon inflow events in freshwater ecosystem of Kurnur Dam, Maharashtra (Patil and More, 2016).

Alkalinity ranged between 86.4-108.4 mg/L throughout study period exhibiting evident seasonal fluctuations with lower alkalinity observed during monsoon months due to dilution from rainwater and comparatively higher values in summer (May-2024) possibly due to increased concentration effects from evaporation. Similar seasonal trends in alkalinity were reported in freshwater ecosystem of Triveni lake in Amravati, Maharashtra (Khan et al., 2012). Dissolved Oxygen remained relatively stable throughout study period (6.72 to 7.84 mg/L). These findings comparable with those reported in Pond of Antargam village, Telangana (Tamlurkar et al., 2023), where dissolved oxygen level ranged between 5 mg/L and 10.41 mg/L. Similarly, during rainy season dissolved oxygen values slightly lower due to dilution by rainwater and increased turbidity which affects and limits photosynthesis due to reduced light penetration. Such trends also evident in our study which indicates a well-balanced aquatic system capable of supporting aquatic life.

Free CO₂ showed moderate fluctuations throughout seasons with lower levels in winter and summer due to early morning uptake by phytoplankton where higher values in monsoon months as photosynthesis gets reduced. Such trend has been also documented in studies like that of Jamwadi Medium Dam, Yavatmal (Wankhade, 2016). Minimum level of chloride during monsoon period due to dilution effects from rainfall and surface runoff which reduces concentration of salts in water. Steady increase in chloride levels after monsoon months as reaching their peak in May-2024. This increase attributed to elevated evaporation rate, reduced inflow and concentration of dissolved ions under higher temperature and stagnant water conditions. These findings align with chloride values trend of increase in summer those reported in study of Vishnupuri dam, Maharashtra (Ubarhande, 2018).

The observed peak in both nitrate and phosphate concentration during post-monsoon period can be attributed to surface runoff, agricultural leaching and organic matter decomposition collectively increase nutrient loading into aquatic system. Similar seasonal enrichment patterns have been reported in freshwater reservoirs, particularly during post-monsoon months when catchment inflows intensity nutrient input. Observed seasonal pattern showed a slight decline in calcium levels during monsoon months (June 2023 to September 2023) due to dilution effects caused by heavy rainfall and runoff followed by gradual increase till May-2024 which can be attributed to higher evaporation rates, reduced inflow and possible enrichment from soil and mineral weathering in surrounding area.

Slight increase in Biological Oxygen Demand during (June-2023 to August-2023) is likely due to runoff carrying decomposable organic matter from surrounding agricultural fields and soil erosion into the dam. Biological Oxygen Demand gradually decreases reaching its low in February may be due to lower microbial activity during colder months, reduced biological decomposition and higher dissolved oxygen. Onwards March-2024, biological oxygen demand rise again slightly in late summer generally associated with rising temperatures that promote microbial metabolism and decomposition of accumulated organic material, reduced dissolved oxygen levels. A comparable trend was reported by

Mahajan and Pokale(2017) in their study of Mohabala Lake, Chandrapur, Maharashtra where biological oxygen demand values ranged between 2.8 mg/L and 5.8 mg/L with higher values during summer.

Overall, the water quality parameters recorded throughout year were within favourable limits for sustaining zooplankton communities. This community plays a pivotal role in development of fish species particularly during spawning season.

References:

1. Aazami, J., Esmaili-sari., Abdoli,A., Sohrabi,H.,Van den Brink,P.J.(2015).Monitoring and assessment of water health quality in the Tajan River, Iran using physicochemical, fish and macroinvertebrates indices. Journal of Environmental Health Science and Engineering, 13(29),1-12.
2. Ajagekar V. V., Nikam K.N., Pawar, C.V. (2011).Physico-chemical Limnology of Chitri Reservoir Ajara, Maharashtra. Nature Environment and Pollution Technology,10 (1),145-146.
3. APHA, (1999). Standard Method for Examination of water and waste water (20th ed.).Washington D.C.:American Public Health Association.
4. APHA, (2017). Standard Method for Examination of water and waste water (23rd ed.).Washington D.C.: American Public Health Association.
5. Ayoola,S.O.,Ajani,E.K.(2009).Seasonal variation in fish distribution and physico-chemical parameters of wetland areas in Oyo State, Nigeria. International Journal Biology and Chemical Science,3(1),107-116.
6. Bassem,S.M. (2020).Water pollution and aquatic biodiversity. Biodiversity Int J. 4(1):10–16.
7. Dirican,S.(2015).Assessment of Water Quality Using Physico-chemical Parameters of Camligoze Dam Lake in Sivas,Turkey. Ecologia,5(1),1-7.
8. Gaikwad, V.D., Padule, A.R., More, R.B.(2021).Study of Physico-Chemical Parameters of Uttarmand reservoir in Patan tehsil, District Satara (M.S) India. International Journal of Creative Research Thoughts,9 (7),188-190.
9. Gurchal,S.S., Bhatkulkar,M.M.(2024). Fish Diversity of Pimpalgaon (Wakhaji) dam of Nagpur District (M.S). International Research Journal of Commerce, Arts and Science,15(12),116-123.
10. Khan, R.M., Jadhav, M.J., Ustad, I.R.(2012). Physicochemical Analysis of Triveni lake Water of Amravati District in (M.S) India. Bioscience Discovery,3(1),64-66.
11. Mahajan,V.S.,Pokale,S.S.(2017).Studies on Physico-chemical analysis of Mohabala lake near Bhadrawati,District-Chandrapur(M.S),India.International Journal of Life Sciences,5(3),438-446.
12. Makode,P.M.(2020).Water quality analysis of Shivan irrigation Dam Near Murtizapur, Dist.Akola. International Research Journal of Science and Engineering,A7,411-416.
13. Patil,S.S.,More,V.R.(2016).Seasonal changes in zooplankton community structure in relation to physico-chemical factors at Kurnur dam

14. Dist.Solapur(MS).International Journal of Fauna and Biological Studies,4(1),86-92.
15. Rathod,S.D.,Walale,M.P.(2022).Study on zooplankton diversity of Yaldali Dam,Parbhani (M.S),India.International Journal of Life Sciences,10(4),375-380.
16. Shinde, S.E., Pathan, T.S., Raut, S.K., Sonawane, D.L.(2011).Studies on the physico-chemical parameters and correlation coefficient of Harsool-Savangi Dam, District Aurangabad, India. Middle-East Journal of Scientific Research, 8 (3),544-554.
17. Tamlurkar, H., Chadrashekhkar,P., Sandhya Rani, B., Umme,T.(2023). Seasonal Variation in Dissolved Oxygen of Pond Water in Antargam Village, Jagtial, Jagtial Dist.Telangana State. Biolife, 11(2),31-33.
18. Ubarhande,S.(2018).Study of Physico-Chemical parameters from Vishnupuri Dam, Nanded, Maharashtra, India.Journal of Emerging Technologies and Innovative Research,(5)12,16-28.
19. Vyas, A., Radadia, B.,Pandya, M., Vyas, K. (2024). Assessment of Physico-Chemical Parameters and Zooplankton Diversity in Anandpur Dam, Junagadh, Gujarat, India. Journal of Advanced Zoology, 45(2),287-298.
20. Wanjari, H.V.,Tanpure,O.G.(2020).Water quality assessment of Adol Dam, Washim (M.S) India. International Journal of Creative Research Thoughts,8(4), 652-659.
21. Wankhade,R,R.(2016). Study of Seasonal Temperature changes and their influence on free Carbon-dioxide, Dissolved Oxygen (DO) and pH in Jamwadi medium Dam, Jamwadi, Dist-Yavatmal, MS, India. Research Journal of Chemical Sciences,6(4),41-43.
22. Wankhade, V.,Manwar,N.,Malu, A.(2012). Evaluation of status of ecosystem of Sawanga (Vithoba) Lake (Malkhed Talav), District Amravati, Maharashtra by assessment of Some Physicochemical Characteristics of water. International Journal of Scientific and Research Publications, 2 (8), 1-10.